

Florida Department of Environmental Protection Office of Ecosystem Projects

Nutrient Trends in the Kissimmee-Okeechobee-Everglades Ecosystem

August 6, 2015
















Kissimmee-Okeechobee-Everglades Ecosystem






Total Nitrogen & Total Phosphorous Trends (2005-2015)







Northern Everglades Total Nitrogen Trends





 Overall TN concentrations have qualitatively decreased throughout the period of record (WY1979 – 2015)¹.

¹ Due to limited data availability no statistical analysis was performed on the entire POR.




Northern Everglades Total Nitrogen Trends







Northern Everglades **Total Phosphorous Trends**




Limited data available


- Overall TP has fluctuated throughout the period of record (WY1979 – 2015).
- In recent years a steady qualitative decline in TP concentration have been observed from WY1999 to recent¹.

¹ Due to limited data availability no statistical analysis was performed on the entire POR.



Northern Everglades Total Phosphorous Trends







Northern Everglades Nutrients Trends

- On-farm BMPs
- Lakeside Ranch STA
- Taylor Creek/Nubbin Slough STAs
- KRR restoration
- Hybrid Wetland Treatment Technology

Magnitude of Change Summary Total Nitrogen

Statistics	Value (mg/L TN /Year)
Mean	-0.004
Minimum	-0.07
Maximum	0.06


Total Phosphorous

Statistics	Value (µg/L TP /Year)
Mean	-1.4
Minimum	-20.9
Maximum	25.1

Period of Record is 2005 to 2015




Everglades Agricultural Area






Everglades Agricultural Area





Everglades Protection Area Total Nitrogen Trends




- Overall declining trend in regional TN geometric mean concentrations for both inflow and interior regions throughout the period of record (WY1979 – 2015).
- More detailed trend analysis can be found in the 2015 and 2016 South Florida Environmental Report (Ch 3A).


¹ Inflow and interior annual geometric mean concentrations are limited to WCA-1,2 and 3.



Everglades Protection Area Total Nitrogen Trends








2015 TN Annual Geometric Mean (mg/L)





Everglades Protection Area Total Phosphorous Trends





- Overall declining trend in regional TP geometric mean concentrations for both inflow and interior regions throughout the period of record (WY1979 – 2015).
- More detailed trend analysis can be found in the 2015 and 2016 South Florida Environmental Report (Ch 3A).


¹ Inflow and interior annual geometric mean concentrations are limited to WCA-1,2 and 3.



Everglades Protection Area Total Phosphorous Trends







2015 TP Annual Geometric Mean (µg/L)





Everglades Protection Area Nutrient Trends

Projects and BMPs

- On farm BMPs
- STAs
 - 2005: All STAs were operating

Magnitude of Change Summary

Total Nitrogen

Statistics	Value (mg/L TN /Year)
Mean	-0.009
Minimum	-0.08
Maximum	0.05


Total Phosphorous

Statistics	Value (µg/L TP /Year)
Mean	-0.26
Minimum	-2.72
Maximum	5.74


Period of Record is 2005 to 2015



Phosphorus Trends in the Everglades Protection Area




- Water quality improvement projects have reduced the total load of phosphorus entering the downstream ecosystem
- Technical analyses of water quality conditions in the Everglades Protection Area show downward trends in total phosphorus levels within the downstream ecosystem
- Analyses inform on-going interagency management and restoration efforts



Julian P, Payne GG, Xue SK (2016) Chapter 3A: Water Quality in the Everglades Protection Areas. 2016 South Florida Environmental Report



Future TP Reductions



Based on the Governor's Funding Plan

Northern:

TP – 26 MT/year

• Eastern:

TP - 63 MT/year

Western:

TP – 9 MT/year

• Southern:

TP – 86 MT/year



Questions?

Questions?

8/6/2015 18